arXiv:hep-ph/0004120v1 13 Apr 2000

hep-ph/0004120
LMU-00-04
April 2000

A Novel Possibility to Determine the CP-violating Phase γ and the B^0_s-$\bar{B}^0_s$ Mixing Parameter y_s at the $\Upsilon(5S)$ Resonance

Zhi-zhong Xing [*]
Sektion Physik, Universität München, Theresienstrasse 37A, 80333 München, Germany

Abstract

We show that a CP-violating phase can be model-independently determined from the time-independent measurement of coherent $B^0_s\bar{B}^0_s$ decays into $(D^{(*)\pm}_s K^{(*)\mp})(D^{(*)\pm}_s K^{(*)\mp})$ states at the $\Upsilon(5S)$ resonance. This phase amounts to $(-\gamma)$ within the standard model, where γ is the well-known angle of the quark mixing unitarity triangle. It is also possible to determine or constrain the B^0_s-$\bar{B}^0_s$ mixing parameter y_s from the same measurement.

---

[*] E-mail: Xing@hep.physik.uni-muenchen.de

Today the weak decays of B_d and B_s mesons are playing crucial roles in the study of flavor mixing and CP violation beyond the neutral kaon system. The best place to produce B^0_d and $\bar{B}^0_d$ events with high statistics and low backgrounds is the $\Upsilon(4S)$ resonance, on which both the symmetric B factory at Cornell and the asymmetric B factories at KEK and SLAC are based. Similarly a wealth of coherent B^0_s and $\bar{B}^0_s$ mesons can be achieved at the $\Upsilon(5S)$ resonance. Recently some interest has been paid to the possibilities to investigate CP violation and probe new physics in weak B_s decays at the $\Upsilon(5S)$ resonance [1, 2], although it remains an open question whether the existing B factories running at the $\Upsilon(4S)$ energy threshold will finally be updated to run at the $\Upsilon(5S)$ energy threshold.

In this paper we point out a novel idea, which works for the coherent decays of $B^0_s\bar{B}^0_s$ pairs into $(D^\pm_s K^\mp)(D^\pm_s K^\mp)$, $(D^{*\pm}_s K^\mp)(D^{*\pm}_s K^\mp)$, $(D^\pm_s K^{*\mp})(D^\pm_s K^{*\mp})$ and $(D^{*\pm}_s K^{*\mp})(D^{*\pm}_s K^{*\mp})$ states at the $\Upsilon(5S)$ resonance, to extract the CP-violating phase

$$\phi \;\equiv\; \arg\left[\frac{q}{p}\cdot\frac{V_{ub}V^*_{cs}}{V^*_{cb}V_{us}}\right] \; , \tag{1}$$

where q/p describes the weak phase of B^0_s-$\bar{B}^0_s$ mixing, and V_{ub}, V_{cs}, V_{cb} and V_{us} are four elements of the Cabibbo-Kobayashi-Maskawa (CKM) quark mixing matrix. Within the standard model $q/p = (V^*_{tb}V_{ts})/(V_{tb}V^*_{ts})$ is an excellent approximation, therefore ϕ amounts to $(-\gamma)$ to a good degree of accuracy, where

$$\gamma \;\equiv\; \arg\left[-\frac{V^*_{ub}V_{ud}}{V^*_{cb}V_{cd}}\right] \tag{2}$$

denotes one inner angle of the well-known CKM unitarity triangle [3]. So far numerous methods have been proposed towards a clean determination of γ in the weak decays of B mesons [4]. The advantages of our present approach are remarkable: (1) it is completely independent of specific models or approximate symmetries in treating the relevant hadronic matrix elements of B transitions; (2) its feasibility does not require any time-dependent measurement of B_s decays, which is quite difficult due to the expected rapid rate of B^0_s-$\bar{B}^0_s$ oscillation; and (3) it remains valid to determine ϕ and able to shed light on γ, even if there exists a kind of yet unknown new physics in B^0_s-$\bar{B}^0_s$ mixing.

The transitions $B^0_s \rightarrow D^{(*)-}_s K^{(*)+}$ and $\bar{B}^0_s \rightarrow D^{(*)-}_s K^{(*)+}$ occur only through the tree-level quark diagrams with the CKM factors $(V^*_{cb}V_{us})$ and $(V_{ub}V^*_{cs})$, respectively (see Fig. 1 for illustration). Their CP-conjugate processes $\bar{B}^0_s \rightarrow D^{(*)+}_s K^{(*)-}$ and $B^0_s \rightarrow D^{(*)+}_s K^{(*)-}$ have the corresponding CKM factors $(V_{cb}V^*_{us})$ and $(V^*_{ub}V_{cs})$. Hence each of the four decay amplitudes involves only a single weak phase and a single strong phase [5]. For the study of CP violation it is convenient to define two rephasing-invariant measurables:

$$\begin{aligned}\lambda_f \;&\equiv\; \frac{q}{p}\cdot\frac{\langle f|\bar{B}^0_s\rangle}{\langle f|B^0_s\rangle} \; , \\ \lambda_{\bar{f}} \;&\equiv\; \frac{q}{p}\cdot\frac{\langle \bar{f}|\bar{B}^0_s\rangle}{\langle \bar{f}|B^0_s\rangle} \; , \end{aligned} \tag{3}$$



Figure 1: Quark diagrams for B_s^0 and $\bar{B}_s^0$ decays into $D_s^- K^+$, $D_s^{*-} K^+$, $D_s^- K^{*+}$ or $D_s^{*-} K^{*+}$.

where f and $\bar{f}$ are charge-conjugate states:

$$
\begin{aligned}
f &= D_s^- K^+ \ , \ D_s^{*-} K^+ \ , \ D_s^- K^{*+} \ , \ D_s^{*-} K^{*+} \ ; \\
\bar{f} &= D_s^+ K^- \ , \ D_s^{*+} K^- \ , \ D_s^+ K^{*-} \ , \ D_s^{*+} K^{*-} \ .
\end{aligned} \tag{4}
$$

Note that $|q/p| = 1$ holds up to an accuracy of $\mathcal{O}(10^{-4})$ in the standard model [6], and it is expected to remain valid up to an accuracy of $\mathcal{O}(10^{-2})$ even if there exists a kind of yet unknown new physics with large CP violation in B_s^0-$\bar{B}_s^0$ mixing [1, 7]. We therefore take $|q/p| \approx 1$ as a good approximation in the subsequent discussions. Then λ_f and $\lambda_{\bar{f}}$ can be explicitly parametrized as [5]

$$
\begin{aligned}
\lambda_f &= \rho \ e^{i(\phi+\delta)} \ , \\
\lambda_{\bar{f}} &= \frac{1}{\rho} \ e^{i(\phi-\delta)} \ ,
\end{aligned} \tag{5}
$$

where ϕ is the overall weak phase defined already in Eq. (1), δ is the relevant strong phase difference, and

$$
\rho \equiv \left| \frac{\langle f|\bar{B}_s^0\rangle}{\langle f|B_s^0\rangle} \right| = \left| \frac{\langle \bar{f}|B_s^0\rangle}{\langle f|B_s^0\rangle} \right| \tag{6}
$$

measures the ratio of two decay amplitudes. Of course both ρ and δ depend upon the specific final state f.

The coherent B_s^0 and $\bar{B}_s^0$ pairs with odd or even charge-conjugation (C) parity can be produced at the $\Upsilon(5S)$ resonance [8]. The joint decay rates of $B_s^0\bar{B}_s^0$ mesons into f_1 and f_2 states, in both $C=-1$ and $C=+1$ cases, have been derived in Refs. [9, 10]. Here we are only interested in the time-independent measurements. The generic formula for the time-integrated rate of a joint $B_s^0\bar{B}_s^0$ decay mode reads [9]

$$
\begin{aligned}
\mathcal{R}(f_1, f_2)_C \ \propto \ & |\langle f_1|B_s^0\rangle|^2 |\langle f_2|B_s^0\rangle|^2 \left[\frac{1+Cy_s^2}{(1-y_s^2)^2} \left(|\xi_C|^2 + |\zeta_C|^2 \right) - \frac{2(1+C)y_s}{(1-y_s^2)^2} \mathrm{Re}\left(\xi_C^* \zeta_C\right) \right. \\
& \left. - \frac{1-Cx_s^2}{(1+x_s^2)^2} \left(|\xi_C|^2 - |\zeta_C|^2 \right) + \frac{2(1+C)x_s}{(1+x_s^2)^2} \mathrm{Im}\left(\xi_C^* \zeta_C\right) \right] \ ,
\end{aligned} \tag{7}
$$

where $x_s \equiv \Delta M/\Gamma$ and $y_s \equiv \Delta\Gamma/(2\Gamma)$ are two dimensionless parameters of B^0_s-$\bar{B}^0_s$ mixing, and

$$\begin{aligned} \xi_C &= \frac{p}{q}\left(1 + C\lambda_{f_1}\lambda_{f_2}\right) \; , \\ \zeta_C &= \frac{p}{q}\left(\lambda_{f_2} + C\lambda_{f_1}\right) \; . \end{aligned} \tag{8}$$

The definition for λ_{f_1} and λ_{f_2} is similar to that for λ_f in Eq. (3). The present experimental bound for x_s is $x_s \geq 14$ at the 95% confidence level [3]. In addition, a detailed calculation based on the standard model yields $y_s \sim \mathcal{O}(10^{-2})$ up to 0.1 [11]. This value will always be reduced, if B^0_s-$\bar{B}^0_s$ mixing receives CP-violating contributions from new physics [1, 12]. Therefore the formula in Eq. (7) can be simplified by neglecting the $\mathcal{O}(y^2_s)$ and $\mathcal{O}(x^{-2}_s)$ terms. Up to the corrections of $\mathcal{O}(10^{-2})$, we arrive for $C = -1$ at

$$\mathcal{R}(f_1, f_2)_- \propto |\langle f_1|B^0_s\rangle|^2 |\langle f_2|B^0_s\rangle|^2 \left[1 + |\lambda_{f_1}|^2 + |\lambda_{f_2}|^2 + |\lambda_{f_1}|^2|\lambda_{f_2}|^2 - 4{\rm Re}\lambda_{f_1}{\rm Re}\lambda_{f_2}\right] \; ; \tag{9}$$

and for $C = +1$ at

$$\begin{aligned} \mathcal{R}(f_1, f_2)_+ \propto\; & |\langle f_1|B^0_s\rangle|^2 |\langle f_2|B^0_s\rangle|^2 \left[1 + |\lambda_{f_1}|^2 + |\lambda_{f_2}|^2 + |\lambda_{f_1}|^2|\lambda_{f_2}|^2 + 4{\rm Re}\lambda_{f_1}{\rm Re}\lambda_{f_2} \right. \\ & \left. -4y_s\left(1 + |\lambda_{f_1}|^2\right){\rm Re}\lambda_{f_2} - 4y_s\left(1 + |\lambda_{f_2}|^2\right){\rm Re}\lambda_{f_1}\right] \; . \end{aligned} \tag{10}$$

If $y_s \sim \mathcal{O}(10^{-3})$ held, the relevant $\mathcal{O}(y_s)$ terms in $\mathcal{R}(f_1, f_2)_+$ could also be neglected. Here and hereafter we treat y_s as a free parameter of $\mathcal{O}(10^{-2})$. It can be seen later on that the formulas in Eqs. (9) and (10) allow one to determine the weak phase ϕ and the mixing parameter y_s, model-independently, from the joint decays of $B^0_s\bar{B}^0_s$ pairs into (f, f), $(\bar{f}, \bar{f})$ and $(f, \bar{f})$ states.

Now taking $(f_1, f_2) = (f, f)$, $(\bar{f}, \bar{f})$ and $(f, \bar{f})$ respectively, we obtain two ratios of the three joint decay rates for the $C = -1$ case:

$$\begin{aligned} R^{(-)}_f &\equiv \frac{\mathcal{R}(f, f)_-}{\mathcal{R}(f, \bar{f})_-} \\ &= \frac{(1+\rho^2)^2 - 4\rho^2\cos^2(\phi+\delta)}{(1+\rho^2)^2 - 4\rho^2\cos(\phi+\delta)\cos(\phi-\delta)} \; , \\ R^{(-)}_{\bar{f}} &\equiv \frac{\mathcal{R}(\bar{f}, \bar{f})_-}{\mathcal{R}(f, \bar{f})_-} \\ &= \frac{(1+\rho^2)^2 - 4\rho^2\cos^2(\phi-\delta)}{(1+\rho^2)^2 - 4\rho^2\cos(\phi+\delta)\cos(\phi-\delta)} \; ; \end{aligned} \tag{11}$$

and another two ratios for the $C = +1$ case:

$$\begin{aligned} R^{(+)}_f &\equiv \frac{\mathcal{R}(f, f)_+}{\mathcal{R}(f, \bar{f})_+} \\ &= \frac{(1+\rho^2)^2 + 4\rho^2\cos^2(\phi+\delta) - 8y_s\rho(1+\rho^2)\cos(\phi+\delta)}{(1+\rho^2)^2 + 4\rho^2\cos(\phi+\delta)\cos(\phi-\delta) - 8y_s\rho(1+\rho^2)\cos\phi\cos\delta} \; , \\ R^{(+)}_{\bar{f}} &\equiv \frac{\mathcal{R}(\bar{f}, \bar{f})_+}{\mathcal{R}(f, \bar{f})_+} \\ &= \frac{(1+\rho^2)^2 + 4\rho^2\cos^2(\phi-\delta) - 8y_s\rho(1+\rho^2)\cos(\phi-\delta)}{(1+\rho^2)^2 + 4\rho^2\cos(\phi+\delta)\cos(\phi-\delta) - 8y_s\rho(1+\rho^2)\cos\phi\cos\delta} \; ; \end{aligned} \tag{12}$$

The four observables $R^{(\pm)}_f$ and $R^{(\pm)}_{\bar{f}}$ depend upon four unknown quantities ρ, ϕ, δ and y_s, therefore the latter can be determined from the former with some discrete ambiguities (only for ϕ and δ). Such ambiguities can be resolved if the coherent $B^0_s\bar{B}^0_s$ decays into four different final states listed in Eq. (4), which involve the universal ϕ and y_s but the different ρ and δ, are taken into account.

Note that $\delta = 0$ or π would leads to

$$R^{(-)}_f \;=\; R^{(-)}_{\bar{f}} \;=\; R^{(+)}_f \;=\; R^{(+)}_{\bar{f}} \;=\; 1 \; . \tag{13}$$

In this case it is impossible to extract the weak phase ϕ from these observables. However, the four strong phases $\delta(D_sK)$, $\delta(D^*_sK)$, $\delta(D_sK^*)$ and $\delta(D^*_sK^*)$ are in general expected to take values different from one another. Thus one can always pin down the magnitude of ϕ from the measurements of $R^{(\pm)}_f$ and $R^{(\pm)}_{\bar{f}}$ for different final states f. A special value of ϕ ($= 0$ or π) could also result in the relationship in Eq. (13), but this possibility should not happen within the standard model, where $\phi \approx \gamma$ with $0 < \gamma < \pi$.

An estimate of ρ in the naive factorization approximation yields $\rho \sim \mathcal{O}(1)$. Therefore the B^0_s-$\bar{B}^0_s$ mixing parameter y_s of $\mathcal{O}(10^{-2})$ may affect the magnitudes of $R^{(+)}_f$ and $R^{(+)}_{\bar{f}}$ significantly, in particular if the $\cos(\phi \pm \delta)$ terms are of $\mathcal{O}(1)$. This provides an interesting opportunity to determine or constrain y_s model-independently and time-independently.

As the branching ratios of B^0_s decays into $D^\mp_s K^\pm$, $D^{*\mp}_s K^\pm$, $D^\mp_s K^{*\pm}$ and $D^{*\mp}_s K^{*\pm}$ states are all at the $\mathcal{O}(10^{-4})$ level [5], to observe signals of the joint $B^0_s\bar{B}^0_s$ decays into these final states needs about 10^8 $B^0_s\bar{B}^0_s$ events at the $\Upsilon(5S)$ resonance. The similar number of $B^0_s\bar{B}^0_s$ pairs is required to apply the idea proposed in Ref. [2] for a determination of γ using the partial rates for CP-tagged B_s decays into $D^{(*)\mp}_s K^{(*)\pm}$ states at the $\Upsilon(5S)$ resonance.

Finally let us give a brief comparison between our new approach to determining the CP-violating phase γ and those presented in Ref. [2] and Ref. [5], which all make use of the B_s decays into $D^{(*)\mp}_s K^{(*)\pm}$ states. The method of Ref. [5] requires the time-dependent measurements, therefore its feasibility relies crucially upon the knowledge of B^0_s-$\bar{B}^0_s$ mixing (i.e., the known values of x_s and y_s). To apply the time-independent method of Ref. [2], one needs the quantitative information on the relevant decay amplitudes of *pure* B^0_s and $\bar{B}^0_s$ mesons, which can be experimentally achieved only after the values of x_s and y_s have been fixed. For our approach, the precise knowledge of B^0_s-$\bar{B}^0_s$ mixing is not a prerequisite. Indeed it is even possible to measure the magnitude of y_s using our method at the $\Upsilon(5S)$ resonance. Within the standard model the three approaches can be complementary to one another for the determination of γ.

In summary, we have proposed a new method to extract the weak phase γ and to determine the B^0_s-$\bar{B}^0_s$ mixing parameter y_s from the coherent $B^0_s\bar{B}^0_s$ decays into $(D^{(*)\mp}_s K^{(*)\pm})(D^{(*)\mp}_s K^{(*)\pm})$ states at the $\Upsilon(5S)$ resonance. It will become realistic and useful, perhaps in the second (or final) round of B-factory experiments, once the present e^+e^- colliders operating at the $\Upsilon(4S)$ resonance are updated to run at the $\Upsilon(5S)$ energy threshold.

The author is grateful to X. Calmet, A.F. Falk, C.D. Lü, and A.A. Petrov for useful communications.

References

[1] Z.Z. Xing, Eur. Phys. J. C **4** (1998) 283.

[2] A.F. Falk and A.A. Petrov, hep-ph/0003321 (to be published).

[3] Particle Data Group, C. Caso *et al.*, Eur. Phys. J. C **3** (1998) 1.

[4] For a review with extensive references, see: The *BABAR* Physics Book, edited by P.F. Harrison and H.R. Quinn, Report No. SLAC-R-504 (1998).

[5] R. Aleksan, I. Dunietz, and B. Kayser, Z. Phys. C **54** (1992) 653.

[6] M. Lusignoli, Z. Phys. C **41** (1989) 645.

[7] See, e.g., A.I. Sanda and Z.Z. Xing, Phys. Rev. D **56** (1997) 6866;
L. Wolfenstein, Phys. Rev. D **57** (1998) 5453;
L. Randall and S. Su, Nucl. Phys. B **540** (1999) 37;
G. Barenboim, J. Bernabéu, and J. Matias, Phys. Rev. D **60** (1999) 016003.

[8] See, e.g., P. Krawczyk, D. London, and H. Steger, Nucl. Phys. B **312** (1989) 1;
J. Lee-Franzini *et al.*, Phys. Rev. Lett. **65** (1990) 2947.

[9] Z.Z. Xing, Phys. Rev. D **55** (1997) 196.

[10] Z.Z. Xing, Phys. Rev. D **53** (1996) 204; Phys. Rev. D **50** (1994) 2957; Phys. Lett. B **463** (1999) 323.

[11] M. Beneke, G. Buchalla, C. Greub, A. Lenz, and U. Nierste, Phys. Lett. B **459** (1999) 631.

[12] Y. Grossman, Phys. Lett. B **380** (1996) 99.